

ERRATUM

Open Access



Erratum to: 'Perioperative use of crystalloids in patients undergoing open radical cystectomy: balanced Ringer's maleate versus a glucose 5 %/potassium-based balanced solution: study protocol for a randomized controlled trial'

Lukas M. Löffel¹, Bettina Kleeb¹, Fiona C. Burkhard² and Patrick Y. Wuethrich^{1*}

Unfortunately, the original version of this article [1] contained a few errors in Table 1. The correct data for Table 1 has now been included below.

Author details

¹Department of Anesthesiology and Pain Medicine, Inselspital, Bern University Hospital, Berne CH-3010, Switzerland. ²Department of Urology, Inselspital, Bern University Hospital, Bern, Switzerland.

Published online: 18 February 2016

Reference

1. Löffel LM, Bettina K, Burkhard FC, Wuethrich PY. Perioperative use of crystalloids in patients undergoing open radical cystectomy: balanced Ringer's maleate versus a glucose 5 %/potassium-based balanced solution: study protocol for a randomized controlled trial. *Trials*. 2014;15:276.

* Correspondence: patrick.wuethrich@insel.ch

¹Department of Anesthesiology and Pain Medicine, Inselspital, Bern University Hospital, Berne CH-3010, Switzerland



Table 1 Electrolyte composition of the 2 different crystalloid solutions

| | Ringerfundin® | G5-K Solution® |
|------------------|---------------|----------------|
| Natrium | 145.0 mmol/l | 50.0 mmol/l |
| Kalium | 4.0 mmol/l | 30.0 mmol/l |
| Magnesium | 1.0 mmol/l | 2.0 mmol/l |
| Calcium | 2.5 mmol/l | 0 mmol/l |
| Chlorid | 127.0 mmol/l | 65 mmol/l |
| Acetat | 24.0 mmol/l | 0 mmol/l |
| Maleat acid | 5.0 mmol/l | 0 mmol/l |
| Lactate | 0 mmol/l | 18 mmol/l |
| HPO ₄ | 0 mmol/l | 8.0 mmol/l |
| Glucose | 0 g/l | 50 g/l |